

Management Practices to Conserve the Fertilizer N Value of Dairy Manure in Vakinankaratra Region, Madagascar

Paulo Salgado¹, Emmanuel Tillard², Bakoly Rarivoarimanana³, Virginie Decruyenaere⁴, Philippe Lecomte⁵

¹UMR SELMET, CIRAD, 110 Antsirabe, Madagascar; ²UMR SELMET, CIRAD, F-97410 Saint-Pierre, La Réunion, France; ³FOFIFA 110 Antsirabe, Madagascar; ⁴CRA-W, Département Productions et Filières, B-5030 Gembloux, Belgium; ⁵UMR SELMET, CIRAD, F-34398 Montpellier, France; paulo.salgado@cirad.fr

Introduction

Madagascar is an island with high agricultural potential, where a great part of the population depends on the primary sector (28% of the GDP; FAOstat, 2011). In Vakinankaratra region, inappropriate agricultural practices forced by an increasing pressure on land led to erosion and decreased soil fertility (Ratsivalaka, 2007). This phenomenon is amplified by the reduced use of mineral fertilizers due to the high prices. But the Vakinankaratra region has the advantage of concentrating large dairy cattle herds, high producers of organic manure. The characterization of manure quality is often addressed by agricultural stakeholders, but it has been little documented in the south countries (Rufino *et al.*, 2006).

Stakeholders involved in agricultural development and research agree on the need to make crop and animal production systems more efficient through smart use of the plant and animal biomass produced on farms, or available in the surrounding environment. Above all, farmers are now aware of the need to restore or maintain the long term fertility of their fields if they want to ensure their own food security or boost their income. In this context, animals are an essential pathway to improve soil fertility by virtue of their ability to collect, convert, use and recycle nutrients.

The objective of this study is to characterize the variation factors that affect manure quality on dairy cattle farms in Vakinankaratra region and to identify practices that increase fertilizer nitrogen value of dairy manure.

Materials and methods

A typology (PCA and hierarchical clustering) of dairy farms in Vakinankaratra region was realised based on data collected on 300 farms. Sixty farms equally sampled in the groups of typology were selected to represent the variability of manure management practices. Three

groups of factors liable to vary manure quality were identified: (i) dairy herd and feed (16 factors), (ii) type of stable and litter management practices (12 factors) and (iii) mode and practices of manure storage (28 factors). The experimental field protocol includes an interview with the herd manager, observations on management practices and samplings. Several hundred samples (65 samples of feeds, 82 faeces, 124 litters and 329 manures) were collected and stored in a freezer. After drying in the oven (24 to 72 hours, at 48 °C) and milling (1 mm), samples were scanned using a near-infrared spectrometer (NIRS) to predict their chemical composition (nitrogen and carbon) and their nutritional and/or fertilizer value (nitrogen).

A Discriminant Correspondence Analysis (Greenacre, 1993) was used to determine management practices specifically associated with the variations of the nitrogen content of organic matter (faeces litter and manure).

Results and discussion

A herd with high number of calves and heifers, adult cows receiving only one meal per day, rice straw as the main forage of the diet and the lack of concentrate feed were linked to the production of faeces with high nitrogen content (low nitrogen digestibility). Jarrige *et al.* (1988) have shown that the use of nitrogen and its digestibility vary according to the age of the animals. The digestive system of young animals is under development and they have a lower digestive capacity, especially for fibrous feeds, compared to that of adult animals. Robison and Sniffen, (1985), showed that increasing meal frequency decreases dietary nitrogen excretion in faeces and urine. Rumen microflora efficiency, and consequently nutrient digestibility, depends on the combined supply of energy and protein. These supplies must be sufficient, equal, simultaneous and continuous throughout the day. Ruminants that receive only one meal per day or a diet based on rice straw and without supply of concentrates have a non-optimal microbial activity, a decrease of the microbial digestion efficiency and therefore a low digestibility of dietary nitrogen.

The highest nitrogen content (1.9 to 2.6% DM) of scrapped litters appears most often on farms with rice straw litters and stables with paved floor. The fiber composition of the plant biomass used as litter plays a role in the absorption of urine during storage and scraping (Lekasi *et al.*, 2003). Nzuma and Murwira (2000) have demonstrated that the addition of straw to the litter allowed reducing losses of urinary nitrogen by ammonia volatilization up to 85%. Compared with bare soils, paved soils avoid that faeces and urine infiltrate into the soil and thereby allows moistening and enriching the straw used in the litter. Plant litter is

easier to decompose thanks to urea in particular (biological action), but it is also degraded thanks to trampling by cattle against the paved floor (mechanical action).

The highest nitrogen content (2.0 to 2.6% DM) of manure come from farms where we observe most often manure storage in ditches, the addition of pig nitrogen-rich manure (> 3% DM), the addition of poultry litter and a manure storage period of less than 90 days. Rufino *et al.* (2006) demonstrated that a manure heap uncovered is much more exposed to losses by leaching and/or evaporation. The air, the rain and the wind amplify the nitrogen volatilization phenomenon in ammoniacal form; the greater the manure surface in contact with air and wind, the greater are the losses.

Conclusion

Adding-value to manure and to other livestock effluents has become essential to maintain soil fertility in Vakinankaratra region. The composition and the fertilizer value of manure are highly variable from one farm to another, and are strongly related to farming systems and to management and storage practices of organic matter. Some practices considered by farmers as capable to improve manure quality, were not recognized as such. They correspond more to a desire, widespread among farmers, of adding-value to all organic matter resources available at the farm level (residues, waste, leaves, etc.) and to recycle everything that is considered "waste" in order to limit input purchase.

These results are important for advising farmers on how to produce quality organic fertilizer and improve fertilization techniques in low-input farming systems. These improvements not only help to increase food self-sufficiency and income levels among farming households, but also to reduce their reliance on outside inputs, the price of which greatly depends on market volatility. Moreover, reducing mineral fertilizer use helps to improve the environmental efficiency of agricultural activities (reduction in greenhouse gas emissions and fossil fuel consumption).

References

- FAOSTAT 2011. Food and Agriculture Organization of the United Nations. [On-line]. [2011/08]. URL : <http://faostat.fao.org>.
- Greenacre MJ. 1993. Correspondence analysis in practice. London. Academic Press.
- Jarrige R. Ed. 1988. Alimentation des bovins, ovins et caprins. INRA, Paris, 476 p.

- Lekasi J.K., Tanner J.C., Kimani S.K. et Harris P.J.C. 2003. Cattle manure quality in Maragua District, Central Kenya: effect of management practices and development of simple methods of assessment. *Agriculture, Ecosystems and Environment* 94:289-298.
- Nzuma J.K. et Murwira H.K. 2000. Improving the Management of Manure in Zimbabwe. *Managing Africa's Soils*, vol. 15. IIED-Drylands Programme, London, UK.
- Ratsivalaka S.R. 2007. Les conditions socio-économiques de la lutte anti-érosive à Madagascar. Actes des JSIRAUF, Hanoi, 6-9 novembre 2007.
- Robinson H. et Sniffen C.J. 1985. Forestomach and whole tract digestibility for lactating dairy cows as influenced by feeding frequency. *J Dairy Sci* 68:857-867.
- Rufino M.C., Rowe E.C., Delve R.J. et Giller K.E. 2006. Nitrogen cycling efficiencies through resource-poor African crop-livestock systems. *Agriculture, Ecosystems and Environment* 112, 261-282.